

## **II. AMENDMENTS TO THE CLAIMS**

Claim 1. (Previously Presented) A film-coated tablet with cyclophosphamide as active compound, comprising in the core cyclophosphamide, one or more fillers, one or more flow regulators, one or more lubricants, and one or more dry binders, wherein as least one of the fillers is lactose monohydrate and wherein the film-coated tablet does not contain preswollen starch.

Claim 2. (Currently Amended) The film-coated tablet as claimed in claim 1, comprising in the core as a filler lactose monohydrate, D-mannitol or [CaHPO<sub>4</sub>] CaHPO<sub>4</sub>, nonpreswollen cornstarch or microfine [cellulose] cellulose, as a flow regulator highly disperse silica and as a lubricant magnesium stearate, stearic acid, glycerol palmitostearate, polyethylene glycol, talc or glycerol monobehenate.

Claim 3. (Previously Presented) The film-coated tablet as claimed in claim 2, where the cores can comprise the auxiliaries either individually or alternatively in any desired mixture.

Claim 4. (Currently Amended) The film-coated tablet as claimed in any one of claims 1 to 3, comprising, per part of cyclophosphamide in the core, lactose monohydrate, microfine cellulose, nonpreswollen cornstarch talc, highly disperse silica and magnesium stearate in the following ratios: lactose monohydrate 0.2-1.5, microfine cellulose 0.2-1.5, nonpreswollen cornstarch talc [0.1 - 1.5] 0.1-1.5, highly disperse silica [0.01 -0.1] 0.01-0.1, and magnesium stearate [0.01 -0.1] 0.01-0.1.

Claim 5. (Currently Amended) The film-coated tablet as claimed in claim 4, comprising, per part of cyclophosphamide in the core, lactose monohydrate, microfine cellulose, nonpreswollen cornstarch talc, highly disperse silica and magnesium stearate in the following ratios: lactose monohydrate 0.5-1, microfine cellulose 0.5-1, nonpreswollen cornstarch talc [0.2-0.7,highly] 0.2-0.7, highly disperse silica 0.01-0.5, and magnesium stearate [0.01 -0.05] 0.01-0.05.

Claim 6. (Currently Amended) The film-coated tablet as claimed in claim 5, comprising, per part of cyclophosphamide in the core, lactose monohydrate, microfine cellulose, nonpreswollen cornstarch talc, highly disperse silica and magnesium stearate in the following [ratios:lactose] ratios: lactose monohydrate 0.73, microfine cellulose 0.74, nonpreswollen cornstarch talc 0.37, highly disperse silica 0.04, and magnesium stearate 0.03.